

Future
Neutrino
Physics
Prospects
Using Massive
Detectors at
DUSEL

Mary Bishai
Brookhaven
National
Laboratory

Neutrino
mixing: A
brief overview

CP violation
and the mass
hierarchy

DUSEL and
LBNE

Summary

Future Neutrino Physics Prospects Using Massive Detectors at DUSEL

ACS 2009, Neutrino Symposium, Aug 16-18, Washington
DC

Mary Bishai
Brookhaven National Laboratory

August 13, 2009

Outline

Future
Neutrino
Physics
Prospects
Using Massive
Detectors at
DUSEL

Mary Bishai
Brookhaven
National
Laboratory

Neutrino
mixing: A
brief overview

CP violation
and the mass
hierarchy

DUSEL and
LBNE

Summary

1 Neutrino mixing: A brief overview

2 CP violation and the mass hierarchy

3 DUSEL and LBNE

4 Summary

Neutrinos have flavors

Future
Neutrino
Physics
Prospects
Using Massive
Detectors at
DUSEL

Mary Bishai
Brookhaven
National
Laboratory

Neutrino
mixing: A
brief overview


CP violation
and the mass
hierarchy

DUSEL and
LBNE


Summary



1962: Leon Lederman, Melvin Schwartz and Jack Steinberger use BNL's Alternating Gradient Synchrotron (AGS) to produce a beam of neutrinos using the decay $\pi^+ \rightarrow \mu^+ \nu_\mu$



The AGS



Making ν 's

Result: 40 neutrino interactions recorded in the detector, 6 of the resultant particles were identified as background and 34 identified as $\mu \Rightarrow \nu_x = \nu_\mu$

The first accelerator neutrino experiment was at the AGS.

Neutrino oscillations

Future
Neutrino
Physics
Prospects
Using Massive
Detectors at
DUSEI

Mary Bishai
Brookhaven
National
Laboratory

Neutrino mixing: A brief overview

1957, 1967: B. Pontecorvo proposes that neutrinos could oscillate:

$$\begin{pmatrix} \nu_a \\ \nu_b \end{pmatrix} = \begin{pmatrix} \cos(\theta) & \sin(\theta) \\ -\sin(\theta) & \cos(\theta) \end{pmatrix} \begin{pmatrix} \nu_1 \\ \nu_2 \end{pmatrix}$$

$$\nu_a(t) = \cos(\theta)\nu_1(t) + \sin(\theta)\nu_2(t)$$


$$\begin{aligned} \mathsf{P}(\nu_a \rightarrow \nu_b) &= |<\nu_b|\nu_a(t)>|^2 \\ &= \sin^2(\theta) \cos^2(\theta) |e^{-iE_2 t} - e^{-iE_1 t}|^2 \end{aligned}$$

$$P(\nu_a \rightarrow \nu_b) = \sin^2 2\theta \sin^2 \frac{1.27 \Delta m_{21}^2 L}{E}$$

where $\Delta m_{21}^2 = (m_2^2 - m_1^2)$ in eV², L (km) and E (GeV).

If flavor eigenstates mix with mass

eigenstates \Rightarrow neutrinos have mass



The Homestake Experiment

Future
Neutrino
Physics
Prospects
Using Massive
Detectors at
DUSEL

Mary Bishai
Brookhaven
National
Laboratory

Neutrino
mixing: A
brief overview

CP violation
and the mass
hierarchy

DUSEL and
LBNE

Summary

1967: Ray Davis from BNL installs a large detector, containing 615 tons of tetrachloroethylene (cleaning fluid), 1.6km underground in Homestake mine, SD.

- 1 $\nu_e^{\text{sun}} + {}^{37}\text{CL} \rightarrow e^- + {}^{37}\text{Ar}$, $\tau({}^{37}\text{Ar}) = 35$ days.
- 2 Number of Ar atoms = number of ν_e^{sun} interactions.



Ray Davis

Results: 1969 - 1993 Measured 2.5 ± 0.2 SNU (1 SNU = 1 neutrino interaction per second for 10^{36} target atoms) while theory predicts 8 SNU. This is a ν_e^{sun} deficit of 69%.

Solar ν_e and atmospheric ν_μ disappearance \Rightarrow
experimental proof of oscillations (R. Wendell's talk next)

Neutrino Mixing: 3 flavours

Future
Neutrino
Physics
Prospects
Using Massive
Detectors at
DUSEL

Mary Bishai
Brookhaven
National
Laboratory

Neutrino
mixing: A
brief overview



CP violation
and the mass
hierarchy

DUSEL and
LBNE

Summary

We know now of 3 flavours of neutrinos: The 3 flavour PMNS mixing matrix was developed in 1962 by Maki-Nakagawa-Sakata based on Pontecorvo's earlier work:

$$\begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \end{pmatrix} = \underbrace{\begin{pmatrix} U_{e1} & U_{e2} & U_{e3} \\ U_{\mu 1} & U_{\mu 2} & U_{\mu 3} \\ U_{\tau 1} & U_{\tau 2} & U_{\tau 3} \end{pmatrix}}_{U_{PMNS}} \begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \end{pmatrix}$$



In the past 10 yrs we have measured most
of the U_{PMNS} parameters

$$U_{PMNS} \sim \begin{pmatrix} 0.8 & 0.5 & < 0.20 ?? \\ 0.4 & 0.6 & 0.7 \\ 0.4 & 0.6 & 0.7 \end{pmatrix}, V_{CKM} \sim \begin{pmatrix} 1 & 0.2 & 0.004 \\ 0.2 & 1 & 0.04 \\ 0.009 & 0.04 & 1 \end{pmatrix}$$

In contrast to CKM, large off diagonal terms:

Measuring neutrino mixing - ν_e oscillations

Future
Neutrino
Physics
Prospects
Using Massive
Detectors at
DUSEL

Mary Bishai
Brookhaven
National
Laboratory


Neutrino
mixing: A
brief overview

CP violation
and the mass
hierarchy

DUSEL and
LBNE

Summary

Solar ν_e disappearance constrained $1 \rightarrow 2$ mixing. Precision from reactor $\bar{\nu}_e$ experiments:



* = future reactor $\bar{\nu}_e$ experiments

Measuring neutrino mixing - ν_μ oscillations

Future
Neutrino
Physics
Prospects
Using Massive
Detectors at
DUSEL


Mary Bishai
Brookhaven
National
Laboratory

Neutrino
mixing: A
brief overview

CP violation
and the mass
hierarchy

DUSEL and
LBNE

Summary



* = future accelerator ν_μ experiments

SuperK atmospheric ν_μ disappearance (L/E 3 → 30,000) probes
2 → 3 oscillations (talks by Yang, Raaf).

Neutrino Matrix Parameterization

Future
Neutrino
Physics
Prospects
Using Massive
Detectors at
DUSEL

Mary Bishai
Brookhaven
National
Laboratory

Neutrino
mixing: A
brief overview


CP violation
and the mass
hierarchy

DUSEL and
LBNE

Summary

$$\underbrace{\begin{pmatrix} 1 & 0 & 0 \\ 0 & c_{23} & s_{23} \\ 0 & -s_{23} & c_{23} \end{pmatrix}}_{\nu_\mu \text{ disappearance}} \underbrace{\begin{pmatrix} c_{13} & 0 & e^{i\delta_{CP}} s_{13} \\ 0 & 1 & 0 \\ -e^{i\delta_{CP}} s_{13} & 0 & c_{13} \end{pmatrix}}_{\nu_\mu \rightarrow \nu_e, \text{ reactor } \bar{\nu}_e \text{ disappear}} \underbrace{\begin{pmatrix} c_{12} & s_{12} & 0 \\ -s_{12} & c_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix}}_{\text{solar } \nu_e, \bar{\nu}_e \text{ disappear}}$$

where $c_{\alpha\beta} = \cos \theta_{\alpha\beta}$ and $s_{\alpha\beta} = \sin \theta_{\alpha\beta}$ and δ_{CP} is the CP phase.



$\sin^2 \theta_{13}$: Amount of ν_e in ν_3

$\tan^2 \theta_{23}$: Ratio of $\frac{\nu_\mu}{\nu_\tau}$ in ν_3

$\tan^2 \theta_{12}$: $\frac{\text{Amount of } \nu_e \text{ in } \nu_2}{\text{Amount of } \nu_e \text{ in } \nu_1}$

WE DONT KNOW: $\sin^2 2\theta_{13}, \delta_{cp}, \text{sign}(\Delta m_{31}^2)$

The mixing matrix - 2009

J. Valle, TAUP '09

Future
Neutrino
Physics
Prospects
Using Massive
Detectors at
DUSEL

Mary Bishai
Brookhaven
National
Laboratory


Neutrino
mixing: A
brief overview

CP violation
and the mass
hierarchy



DUSEL and
LBNE

Summary

Maltoni et al, NJP 6 (2004) 122




Schwetz et al, NJP 10 (2008) 113011



**Homestake, SAGE+
GALLEX/GNO,
Super-K, SNO
Borexino**

KamLAND (180 Km)

Valle@TAUP09



... Super-K

**K2K (250 Km)
MINOS (735 Km)**

θ_{13} by 2016

M. Mezzetto, arXiv:0905.2842

Future
Neutrino
Physics
Prospects
Using Massive
Detectors at
DUSEL


Mary Bishai
Brookhaven
National
Laboratory

Neutrino
mixing: A
brief overview

CP violation
and the mass
hierarchy

DUSEL and
LBNE

Summary



CP Violation, Mass Hierarchy and $\nu_\mu \rightarrow \nu_e$

Future
Neutrino
Physics
Prospects
Using Massive
Detectors at
DUSEL

Mary Bishai
Brookhaven
National
Laboratory

Neutrino
mixing: A
brief overview


CP violation
and the mass
hierarchy

DUSEL and
LBNE

Summary

*Appearance probabilities of $\nu_\mu \rightarrow \nu_e$ for different values of the CP phase.
A CP phase $\neq 0, \pi$ implies CP is violated in the lepton sector.*

Normal Hierarchy



CP effects largest $E_\nu < 3$ GeV.

CP Violation, Mass Hierarchy and $\nu_\mu \rightarrow \nu_e$

Future
Neutrino
Physics
Prospects
Using Massive
Detectors at
DUSEL

Mary Bishai
Brookhaven
National
Laboratory

Neutrino
mixing: A
brief overview



CP violation
and the mass
hierarchy

DUSEL and
LBNE

Summary

*Appearance probabilities of $\nu_\mu \rightarrow \nu_e$ for different values of the CP phase.
A CP phase $\neq 0, \pi$ implies CP is violated in the lepton sector.*

Reversed Hierarchy



Matter effects large $E_\nu > 1.5$ GeV.

Strategies for Measuring δ_{cp} and the Mass Hierarchy

Future Neutrino Physics Prospects Using Massive Detectors at DUSEL

Mary Bishai
Brookhaven National Laboratory

Neutrino mixing: A brief overview

CP violation and the mass hierarchy


DUSEL and LBNE

Summary


Long baseline accelerator $\nu_\mu \rightarrow \nu_e$ appearance. 300 kT. MW. yr:

- $\sin^2 2\theta_{13} = 0.02, \delta_{\text{cp}} = 0, \text{normal hierarchy}$
- $\sin^2 2\theta_{13} = 0.02, \delta_{\text{cp}} = \pi, \text{normal hierarchy}$
- $\sin^2 2\theta_{13} = 0.02, \delta_{\text{cp}} = -\pi/2, \text{reverse hierarchy}$

NuMI LE at 810 km, 15 mrad off-axis



WBLE 60 GeV at 1300km, 0° off-axis



Off axis NuMI beam at 810 km

A wide-band beam at 1300km

Wide-band beam spectral information = resolves degeneracies

Physics sensitivity vs baseline

Future
Neutrino
Physics
Prospects
Using Massive
Detectors at
DUSEL

Mary Bishai
Brookhaven
National
Laboratory


Neutrino
mixing: A
brief overview

CP violation
and the mass
hierarchy

DUSEL and
LBNE

Summary

Using a broad-band beam with a peak interaction rate at 2 GeV,
 $\text{FWHM}=3 \text{ GeV}$, a parameterized water Cerenkov detector and
exposure of $5 \text{ MW.yr} (\nu) + 10 \text{ MW.yr} (\bar{\nu})$ (V. Barger *et al.*. Phys. Rev. D 74,
073004 2006):



Minimum value of $\sin^2(2\theta_{13})$ for which the sensitivity is $> 3\sigma$
for (best, 50%, worst) of δ_{cp} values

Longer baselines = larger mass effects

Best sensitivity is for baselines 1200 - 2500km

Deep Underground Science and Engineering Laboratory

Future
Neutrino
Physics
Prospects
Using Massive
Detectors at
DUSEL

Mary Bishai
Brookhaven
National
Laboratory


Neutrino
mixing: A
brief overview

CP violation
and the mass
hierarchy

DUSEL and
LBNE

Summary

July 10, 2007: the National Science Foundation (NSF) selected the University California-Berkeley to produce a technical design for DUSEL at Homestake Mine, SD



DUSEL Timeline

J. Dehmer HEPAP Feb, 2009

DUSEL Working Timeline

- July '08: Internal project review of facility & infrastructure.
- January '09: NSF Project Review #1.
- January '10: NSF Project Review #2.
- December '10: NSF Preliminary Design Review (PDR).
 - Project readiness, plan will be assessed at this milestone.
- Spring '11: Presentation of DUSEL MREFC package to NSB.
- FY13: Earliest construction funding (MREFC) start, if approved.

Planning with potential partners (DOE, international, etc.)
being integrated into above schedule.

Future
Neutrino
Physics
Prospects
Using Massive
Detectors at
DUSEL

Mary Bishai
Brookhaven
National
Laboratory

Neutrino
mixing: A
brief overview

CP violation
and the mass
hierarchy

DUSEL and
LBNE

Summary

The Fermilab NuMI Beamlne

Future
Neutrino
Physics
Prospects
Using Massive
Detectors at
DUSEL




Mary Bishai
Brookhaven
National
Laboratory

Neutrino
mixing: A
brief overview

CP violation
and the mass
hierarchy

DUSEL and
LBNE

Summary



Fermilab Neutrino Beams: Future

Future
Neutrino
Physics
Prospects
Using Massive
Detectors at
DUSEL

Mary Bishai
Brookhaven
National
Laboratory

Neutrino
mixing: A
brief overview


CP violation
and the mass
hierarchy

DUSEL and
LBNE

Summary

The NuMI beamline currently uses 300kW sustained power from the Main Injector.

NuMI is the most powerful neutrino beamline in operation today.
Planned upgrades to NuMI and the Main Injector → 700 kW by 2012.



The proposed Project X at FNAL can produce 2MW

The Long Baseline Neutrino Experiment

Future
Neutrino
Physics
Prospects
Using Massive
Detectors at
DUSEL

Mary Bishai
Brookhaven
National
Laboratory


Neutrino
mixing: A
brief overview

CP violation
and the mass
hierarchy

DUSEL and
LBNE

Summary

A Long Baseline Neutrino Experiment (LBNE) from Fermilab to megaton scale detectors at Homestake is now being designed. CDR late 2010.



The LBNE Collaboration

Future
Neutrino
Physics
Prospects
Using Massive
Detectors at
DUSEL

Mary Bishai
Brookhaven
National
Laboratory

Neutrino
mixing: A
brief overview

CP violation
and the mass
hierarchy

DUSEL and
LBNE

Summary



Collaboration meeting 2/26-2/28, 2009 at UC Davis, CA

DUSEL Detectors: Water Cerenkov

Future
Neutrino
Physics
Prospects
Using Massive
Detectors at
DUSEL

Mary Bishai
Brookhaven
National
Laboratory


Neutrino
mixing: A
brief overview

CP violation
and the mass
hierarchy


DUSEL and
LBNE

Summary

SuperKamiokande : 50kT



DUSEL WCe Module : 120 kT



3 100kT (fiducial) modules, $\approx 55\text{m}$ diameter, $\approx 60\text{m}$ height, 60K 10" PMTs/module (25% coverage)

Known technology 2 – 3× SuperK

Higher backgrounds, low efficiency

DUSEL Detectors: Liquid Argon TPC

Future
Neutrino
Physics
Prospects
Using Massive
Detectors at
DUSEL


Mary Bishai
Brookhaven
National
Laboratory

Neutrino
mixing: A
brief overview

CP violation
and the mass
hierarchy

DUSEL and
LBNE

Summary



Small 175 litre prototype in the NuMI beam. Hand scanning and prelim automated MC studies (Tufts U., Yale)

High efficiency and purity

Requires 100× scale-up - unproven.

LBNE/DUSEL spectra

Future
Neutrino
Physics
Prospects
Using Massive
Detectors at
DUSEL

Mary Bishai
Brookhaven
National
Laboratory



Neutrino
mixing: A
brief overview

CP violation
and the mass
hierarchy

DUSEL and
LBNE

Summary

A preliminary on-axis wide-band beam for LBNE based on the NuMI focusing system has been developed. Water Cerenkov response is based on the SuperK MC. LAr is modeled as a near-perfect detector.



Measurements of δ_{cp} in LBNE

Mark Dierckxsens

Future
Neutrino
Physics
Prospects
Using Massive
Detectors at
DUSEL

Mary Bishai
Brookhaven
National
Laboratory

Neutrino
mixing: A
brief overview


CP violation
and the mass
hierarchy

DUSEL and
LBNE


Summary

(θ_{13} , δ_{cp}) Measurement

1 MW



2.3 MW



DUSEL Sensitivities

WCe, 2.3MW beam, 3 yrs ν + 3 yrs $\bar{\nu}$

Future
Neutrino
Physics
Prospects
Using Massive
Detectors at
DUSEL

Mary Bishai
Brookhaven
National
Laboratory


Neutrino
mixing: A
brief overview

CP violation
and the mass
hierarchy


DUSEL and
LBNE

Summary


$\theta_{13} @ 3,5 \sigma$



Mass hierarchy @ 3,5 σ



CP violation @ 3,5 σ



Summary of sensitivities

The smallest value of $\sin^2 2\theta_{13} @ 3\sigma$:

$$\theta_{13} \neq 0 \quad \text{sign}(\Delta m^2) \quad \text{CPV}$$

0.004	0.014	0.012
$\text{all } \delta_{cp}$	$50\% \delta_{cp}$	

Mark Dierckxens, APS 09

Summary and Conclusions

Future
Neutrino
Physics
Prospects
Using Massive
Detectors at
DUSEL

Mary Bishai
Brookhaven
National
Laboratory

Neutrino
mixing: A
brief overview

CP violation
and the mass
hierarchy

DUSEL and
LBNE

Summary

3 flavor neutrino mixing is now well established, but STRANGE. Large off diagonal terms and almost maximal mixing.

BUT- what we dont know is even more important:

- How small is $\sin^2 2\theta_{13}$? Is it close to the current limit (0.1) or is it very small? Is it 0?
- Is there CP violation (and LFV) in the lepton sector?
- What is the mass hierarchy?
- Are there only 3 generations of leptons?

The LBNE project will determine if $\sin^2 2\theta_{13} > 0.005$ and can measure δ_{cp} and the mass hierarchy for $\sin^2 2\theta_{13} \geq 0.01$.



**There is a goldmine of scientific
discoveries at the end of this rainbow**